arXiv:math/0308002v1 [math.GT] 1 Aug 2003

SOME LIMITS OF THE COLORED JONES POLYNOMIALS OF THE FIGURE-EIGHT KNOT

HITOSHI MURAKAMI

Abstract. We will study the asymptotic behaviors of the colored Jones polynomials of the figure-eight knot. In particular we will show that for certain limits we obtain the volumes of the cone manifolds with singularities along the knot.

1. Introduction

Let $J_N(K;t)$ be the colored Jones polynomial of a knot K associated with the N-dimensional irreducible representation of the Lie algebra $\mathfrak{sl}(2;\mathbb{C})$, normalized so that $J_N(\text{unknot};t)=1$. If we evaluate it at the N-th root of unity $\exp(2\pi\sqrt{-1}/N)$, then its absolute value coincides with Kashaev's invariant [2, 3, 9]. It was conjectured by R. Kashaev [3] that the growth rate of his invariant for large N determines the hyperbolic volume of a hyperbolic knot. His conjecture was generalized by J. Murakami and the author to the following Volume Conjecture:

Conjecture 1.1 (Volume Conjecture)**.** *Let K be any knot. Then*

$$2\pi\lim_{N\to\infty}\frac{\log\left|J_N\left(K;\exp(2\pi\sqrt{-1}/N)\right)\right|}{N}=v_3\|S^3\setminus K\|,$$

where v_3 is the volume of the ideal regular hyperbolic tetrahedron and $\|M\|$ is the Gromov norm (or simplicial volume). In particular if K is hyperbolic, then the right hand side coincides with the volume of the knot complement $\operatorname{Vol}(S^3\setminus K)$.

See [4, 14, 12, 13, 15, 16, 10, 5, 7, 6, 8] for related topics.

In this paper we will study limits of the colored Jones polynomials of the figure-eight knot evaluated at $\exp(2\pi r\sqrt{-1}/N)$ for a fixed number r. Put $\Lambda(z):=-\int_0^z\log|2\sin x|dx$, the Lobachevski function, and $\theta(r):=\arccos\left(\cos(2\pi r)-1/2\right)$ with $0\le\arccos(x)\le\pi$.

Theorem 1.2. *Let r be a real number satisfying $5/6<r<7/6$. Then*

$$2\pi\limsup_{N\to\infty}\frac{\log\left|J_N\left(E;\exp(2\pi r\sqrt{-1}/N)\right)\right|}{N}=\frac{2\Lambda\bigl(\pi r+\theta(r)/2\bigr)-2\Lambda\bigl(\pi r-\theta(r)/2\bigr)}{r}.$$

Moreover if r is irrational or $r=1$, then

$$2\pi\lim_{N\to\infty}\frac{\log\left|J_N\left(E;\exp(2\pi r\sqrt{-1}/N)\right)\right|}{N}=\frac{2\Lambda\bigl(\pi r+\theta(r)/2\bigr)-2\Lambda\bigl(\pi r-\theta(r)/2\bigr)}{r},$$

and if $r\neq1$ and rational, then

$$2\pi\liminf_{N\to\infty}\frac{\log\left|J_N\left(E;\exp(2\pi r\sqrt{-1}/N)\right)\right|}{N}=0$$

Date: February 1, 2008.

2000 *Mathematics Subject Classification.* Primary 57M27; Secondary 57M25.

Key words and phrases. colored Jones polynomial, knot, figure-eight knot, volume conjecture, Lobachevski function, cone-manifold.

This research is partially supported by Grant-in-Aid for Scientific Research (B).

Remark 1.3. The case where $r = 1$ is due to R. Kashaev [3] and T. Ekholm [5], and the following proof is similar to Ekholm's.

Remark 1.4. The value $2\Lambda\bigl(\pi r+\theta(r)/2\bigr)-2\Lambda\bigl(\pi r-\theta(r)/2\bigr)$ coincides with the volume of the cone-manifold with underlying space S^3, the singularity the figure-eight knot, and the cone-angle $2\pi|1-r|$ [11, 1], which was informed by S. Gukov, A. Mednykh, and A. Vesnin.

Remark 1.5. Some results in [8] were erroneously stated. The author did not consider the case where r is rational.

We also calculate the limits for some other cases.

2. PRELIMINARIES

Let E denote the figure-eight knot 4_1. Due to K. Habiro and T. Le, the following formula is known.

$$J_N(E;t)=\sum_{k=0}^{N-1}\prod_{j=1}^{k}\left(t^{(N+j)/2}-t^{-(N+j)/2}\right)\left(t^{(N-j)/2}-t^{-(N-j)/2}\right). \tag{2.1}$$

Put $t=\exp(2\pi r\sqrt{-1}/N)$. Since

$$t^{(N+j)/2}-t^{-(N+j)/2}=2\sqrt{-1}\sin(\pi r(N+j)/N)$$

and

$$t^{(N-j)/2}-t^{-(N-j)/2}=2\sqrt{-1}\sin(\pi r(N-j)/N),$$

we have

$$\left(t^{(N+j)/2}-t^{-(N+j)/2}\right)\left(t^{(N-j)/2}-t^{-(N-j)/2}\right) \\ =4\sin(\pi rj/N+\pi r)\sin(\pi rj/N-\pi r).$$

Put

$$\begin{aligned} g(j) :&= 4\sin(\pi rj/N+\pi r)\sin(\pi rj/N-\pi r) \\ &= 2\cos(2\pi r)-2\cos(2\pi rj/N) \end{aligned}$$

and

$$f(k) := \prod_{j=1}^{k} g(j)$$

so that $J_N(E;\exp(2\pi r\sqrt{-1}/N))=\sum_{k=0}^{N-1}f(k)$.

3. THE CASE WHERE r IS IRRATIONAL AND $5/6<r<1$ OR $r=1$.

In this section we prove Theorem 1.2 in the case where $5/6<r<1$ and r is irrational, or $r=1$.

We put

$$B:=\frac{N(1-r)}{r},\quad C:=\frac{N\theta(r)}{2\pi r},\quad\text{and}\quad D:=\frac{N(2\pi-\theta(r))}{2\pi r}$$

with $\theta(r):=\arccos\left(\cos(2\pi r)-1/2\right)$. Here arccos takes its value between 0 and π. Note that $0\le B<C<D<1$ ($B=0$ only if $r=1$), that $g(B)=0$ and $g(C)=g(D)=1$, and that B is not an integer for any N since r is irrational. Therefore $g(j)\neq 0$ for any integer $0<j<N$ in this case.

Since we have

$$\begin{aligned} -1 < g(j) < 0 \quad &\text{for} \quad 0 < j < B, \\ 0 < g(j) < 1 \quad &\text{for} \quad B < j < C, \\ g(j) > 1 \quad &\text{for} \quad C < j < D, \quad \text{and} \\ 0 < g(j) < 1 \quad &\text{for} \quad D < j < N, \end{aligned}$$

Table of $g(j)$ regarding j as a continuous parameter

j	0	$\cdots$	B	$\cdots$	C	$\cdots$	D	$\cdots$	N
$g(j)$	$-1 < g(0) < 0$	$\nearrow$	0	$\nearrow$	1	> 1	1	$\searrow$	0

we see

(1) If $j < B$ then the signs of $f(j)$ alternate, that is, $f(j-1)f(j) < 0$, and if $j > B$ then the signs of $f(j)$ are constant, and
(2) $1 = |f(0)| > |f(1)| > \cdots > |f(\lfloor C \rfloor - 1)|$ and $|f(\lfloor C \rfloor)| < \cdots < |f(\lfloor D \rfloor)|$, where $\lfloor x \rfloor$ is the greatest integer that does not exceed x.

Put $F_N := |f(\lfloor D \rfloor)|$, which is the maximum of $\{|f(j)|\}$ for $0 \le j \le N-1$.

We will show the following inequality.

Claim 3.1.

$$F_N - 1 \le \left| J_N(E; \exp(2\pi r \sqrt{-1}/N)) \right| \le N F_N$$

Proof. The last inequality follows since $|f(j)| \le F_N$ and $J_N(E; \exp(2\pi r\sqrt{-1}/N)) = \sum_{j=0}^{N-1} f(j)$.

To prove the first inequality, first we consider the case where $\lfloor B \rfloor$ is odd. In this case since $f(0) = 1$, $f(2j-1) + f(2j) < 0$ for $2j < \lfloor B \rfloor$, and $f(j) < 0$ for $j \ge \lfloor B \rfloor$, we have

$$\begin{aligned} &\left| J_N(E; \exp(2\pi r\sqrt{-1}/N)) \right| \\ &= \big| f(0) + \{f(1) + f(2)\} + \{f(3) + f(4)\} + \cdots + \{f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)\} \\ &\quad + f(\lfloor B \rfloor) + f(\lfloor B \rfloor + 1) + \cdots + f(N-1) \big| \\ &= -1 + |f(1) + f(2)| + |f(3) + f(4)| + \cdots + |f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)| \\ &\quad + |f(\lfloor B \rfloor)| + |f(\lfloor B \rfloor + 1)| + \cdots + |f(\lfloor D \rfloor)| + \cdots + |f(N-1)| \\ &> F_N - 1 \end{aligned}$$

and the first equality follows.

Next we consider the case where $\lfloor B \rfloor$ is even. In this case since $f(2j) + f(2j+1) > 0$ for $2j + 1 < \lfloor B \rfloor$, and $f(j) > 0$ for $j \ge \lfloor B \rfloor$, we have

$$\begin{aligned} &\left| J_N(E; \exp(2\pi r\sqrt{-1}/N)) \right| \\ &= \big| \{f(0) + f(1)\} + \{f(2) + f(3)\} + \cdots + \{f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)\} \\ &\quad + f(\lfloor B \rfloor) + f(\lfloor B \rfloor + 1) + \cdots + f(N-1) \big| \\ &= |f(0) + f(1)| + |f(2) + f(3)| + \cdots + |f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)| \\ &\quad + |f(\lfloor B \rfloor)| + |f(\lfloor B \rfloor + 1)| + \cdots + |f(\lfloor D \rfloor)| + \cdots + |f(N-1)| \\ &> F_N. \end{aligned}$$

and the first equality also follows. □

Now we study the asymptotic behavior of F_N.

Claim 3.2.

$$\lim_{N \to \infty} \frac{\log F_N}{N} = \frac{1}{\pi r} \left\{ \Lambda\left(\pi r + \theta(r)/2\right) - \Lambda\left(\pi r - \theta(r)/2\right) \right\}.$$

Proof. Since

$$
\begin{aligned}
F_N = |f(\lfloor D \rfloor)| &= \prod_{j=1}^{\lfloor D \rfloor} |g(j)| \\
&= \prod_{j=1}^{\lfloor D \rfloor} |2\sin(\pi r j/N + \pi r)|\, |2\sin(\pi r j/N - \pi r)|\,,
\end{aligned}
$$

we have

$$
\begin{aligned}
&\lim_{N\to\infty} \frac{\log F_N}{N} \\
&= \lim_{N\to\infty} \frac{1}{N} \sum_{j=1}^{\lfloor D \rfloor} \{\log|2\sin(\pi r j/N + \pi r)| + \log|2\sin(\pi r j/N - \pi r)|\} \\
&= \frac{1}{\pi r} \int_{\pi r}^{\pi - \theta(r)/2 + \pi r} \log|2\sin x|\, d\,x + \frac{1}{\pi r} \int_{-\pi r}^{\pi - \theta(r)/2 - \pi r} \log|2\sin x|\, d\,x \\
&= \frac{1}{\pi r} \{-\Lambda(\pi - \theta(r)/2 + \pi r) + \Lambda(\pi r) - \Lambda(\pi - \theta(r)/2 - \pi r) + \Lambda(-\pi r)\} \\
&= \frac{1}{\pi r} \{\Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2)\}\,.
\end{aligned}
$$

Here we use the formulas $-\Lambda(-x) = \Lambda(x + \pi) = \Lambda(x)$. □

Since $\Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2) > 0$ (see Appendix), F_N grows exponentially and so $\lim_{N\to\infty} \log(F_N - 1)/N = \lim_{N\to\infty} \log F_N/N$.

Now since $\lim_{N\to\infty} \log N/N = 0$, we have from Claim 3.1

$$
\begin{aligned}
\lim_{N\to\infty} \frac{\log|J_N(E; \exp(2\pi r\sqrt{-1}/N))|}{N} &= \lim_{N\to\infty} \frac{\log F_N}{N} \\
&= \frac{1}{\pi r} \{\Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2)\}\,.
\end{aligned}
$$

This completes proof of the theorem in the case where r is irrational with $5/6 < r < 1$ or $r = 1$.

4. THE CASE WHERE r IS IRRATIONAL AND $1 < r < 7/6$

We put

$$
B := \frac{N(r-1)}{r}, \quad C := \frac{N\theta(r)}{2\pi r}, \quad D := \frac{N(2\pi - \theta(r))}{2\pi r}, \quad \text{and} \quad B' := \frac{N(2-r)}{r}
$$

Note that $0 < B < C < D < 1$, that $g(B) = g(B') = 0$ and $g(C) = g(D) = 1$, and that B and B' are not integers for any N since r is irrational. Therefore $g(j) \neq 0$ for any integer $0 < j < N$ in this case.

Since we have

$$
\begin{aligned}
-1 < g(j) < 0 &\quad \text{for} \quad 0 < j < B, \\
0 < g(j) < 1 &\quad \text{for} \quad B < j < C, \\
g(j) > 1 &\quad \text{for} \quad C < j < D, \\
0 < g(j) < 1 &\quad \text{for} \quad D < j < B', \quad \text{and} \\
-1 < g(j) < 0 &\quad \text{for} \quad B' < j < N,
\end{aligned}
$$

the proof is similar to the case where $5/6 < r < 1$. We put $F_N := |f(\lfloor D \rfloor)|$.

Table of $g(j)$ regarding j as a continuous parameter

j	0	$\cdots$	B	$\cdots$	C	$\cdots$	D	$\cdots$	B'	$\cdots$	N
$g(j)$	$-1 < g(0) < 0$	$\nearrow$	0	$\nearrow$	1	$g(j) > 1$	1	$\searrow$	0	$-1 < g(j) < 0$	0

If $\lfloor B \rfloor$ is odd, we have

$$\begin{aligned}
&\left|J_N(E;\exp(2\pi r\sqrt{-1}/N))\right| \\
&\quad = \bigl|f(0) + \{f(1)+f(2)\} + \{f(3)+f(4)\} + \cdots + \{f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)\} \\
&\qquad + f(\lfloor B \rfloor) + f(\lfloor B \rfloor + 1) + \cdots + f(\lfloor B' \rfloor - 1) \\
&\qquad + f(\lfloor B' \rfloor) + f(\lfloor B' \rfloor + 1) + \cdots + f(N-1)\bigr| \\
&\quad = -1 + |f(1)+f(2)| + |f(3)+f(4)| + \cdots + |f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)| \\
&\qquad + |f(\lfloor B \rfloor)| + |f(\lfloor B \rfloor + 1)| + \cdots + |f(\lfloor D \rfloor)| + \cdots + |f(\lfloor B' \rfloor - 1)| \\
&\qquad + \begin{cases} |f(\lfloor B' \rfloor) + f(\lfloor B' \rfloor + 1)| + \cdots + |f(N-2) - f(N-1)| \\ \hfill \text{if } N - \lfloor B' \rfloor \text{ is even} \\ |f(\lfloor B' \rfloor) + f(\lfloor B' \rfloor + 1)| + \cdots + |f(N-3) - f(N-2)| + |f(N-1)| \\ \hfill \text{if } N - \lfloor B' \rfloor \text{ is odd} \end{cases} \\
&\quad > F_N - 1
\end{aligned}$$

If $\lfloor B \rfloor$ is even,

$$\begin{aligned}
&\left|J_N(E;\exp(2\pi r\sqrt{-1}/N))\right| \\
&\quad = \bigl|\{f(0)+f(1)\} + \{f(2)+f(3)\} + \cdots + \{f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)\} \\
&\qquad + f(\lfloor B \rfloor) + f(\lfloor B \rfloor + 1) + \cdots + f(\lfloor B' \rfloor - 1) \\
&\qquad + f(\lfloor B' \rfloor) + |f(\lfloor B' \rfloor + 1)| + \cdots + |f(N-1)|\bigr| \\
&\quad = |f(0)+f(1)| + |f(2)+f(3)| + \cdots + |f(\lfloor B \rfloor - 2) + f(\lfloor B \rfloor - 1)| \\
&\qquad + |f(\lfloor B \rfloor)| + |f(\lfloor B \rfloor + 1)| + \cdots + |f(\lfloor D \rfloor)| + \cdots + |f(\lfloor B' \rfloor - 1)| \\
&\qquad + \begin{cases} |f(\lfloor B' \rfloor) + f(\lfloor B' \rfloor + 1)| + \cdots + |f(N-2) - f(N-1)| \\ \hfill \text{if } N - \lfloor B' \rfloor \text{ is even} \\ |f(\lfloor B' \rfloor) + f(\lfloor B' \rfloor + 1)| + \cdots + |f(N-3) - f(N-2)| + |f(N-1)| \\ \hfill \text{if } N - \lfloor B' \rfloor \text{ is odd} \end{cases} \\
&\quad > F_N.
\end{aligned}$$

So we have $F_N - 1 < |J_N(E;\exp(2\pi r\sqrt{-1}/N))| < NF_N$, and the result follows, completing the proof of Theorem 1.2.

As a corollary we have

Corollary 4.1.

$$\lim_{r\to 1}\left\{\limsup_{N\to\infty}\frac{\log\left|J_N(E;\exp(2\pi r\sqrt{-1}))\right|}{N}\right\} = \limsup_{N\to\infty}\frac{\log\left|J_N(E;\exp(2\pi\sqrt{-1}))\right|}{N}$$

5. The case where r is rational and $|1-r| < 1/6$.

Put $r := q/p$ with coprime integers p and q.

Suppose first that N is not a multiple of q. Then B and B' are not integers and $g(j) \neq 0$ for any j. So in this case a similar argument for the case where r is irrational applies and we have

$$\frac{\log|J_{N'}(E;\exp(2\pi r\sqrt{-1}/N))|}{N'} \to \frac{1}{\pi r}\left\{\Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2)\right\}$$

for the subsequence $\{N'\}$ of natural numbers which are not multiples of q.

Next suppose that N is a multiple of q, say $N = nq$. Then B is an integer since $B = n(p-q)$ ($n(q-p)$ respectively) if $5/6 < r < 1$ ($1 < r < 7/6$ respectively).

Therefore $-1 < g(j) < 0$ for $0 < j < B$ and $g(j) = 0$ for $j \geq B$ and so $|f(j)| < 1$ for $0 < j < B$ and $f(j) = 0$ for $j \geq B$. Thus we have

$$\begin{aligned}\left|J_N(E;\exp(2\pi r\sqrt{-1}/N))\right| = \left|\sum_{k=0}^{N-1} f(k)\right| = \left|\sum_{k=0}^{B-1} f(k)\right| \leq \sum_{k=0}^{B-1} |f(k)| \\ < B = N|1-1/r|.\end{aligned}$$

Moreover since

$$\begin{aligned}\left|\sum_{k=0}^{B-1} f(k)\right| &= \begin{cases}\{1+f(1)\}+\{f(2)+f(3)\}+\cdots+\{f(\lfloor B\rfloor-2)+f(\lfloor B\rfloor-1)\} \\ \qquad\qquad\text{if } \lfloor B\rfloor \text{ is even} \\ \{1+f(1)\}+\cdots+\{f(\lfloor B\rfloor-3)+f(\lfloor B\rfloor-2)\}+f(\lfloor B\rfloor-1) \\ \qquad\qquad\text{if } \lfloor B\rfloor \text{ is odd}\end{cases} \\ &> 1+f(1) = 1+2\cos(2\pi r)-2\cos(2\pi r/N) \\ &> 2-2\cos(2\pi r/N).\end{aligned}$$

Therefore

$$\begin{aligned}|J_N(E;\exp(2\pi r\sqrt{-1}/N))| > 2-2\cos(2\pi r/N) > 2-2\left(1-\frac{(2\pi r/N)^6}{6!}\right) \\ = (2\pi r)^6/360N^6.\end{aligned}$$

So we finally have

$$\frac{\log\left((2\pi r)^6/360N^6\right)}{N} < \frac{\log|J_N(E;\exp(2\pi r\sqrt{-1}/N))|}{N} < \frac{\log(N|1-1/r|)}{N}$$

and so $\log|J_N(E;\exp(2\pi r\sqrt{-1}/N))|/N$ can be arbitrarily small in this case. Therefore

$$\frac{\log|J_{N''}(E;\exp(2\pi r\sqrt{-1}/N))|}{N''} \to 0$$

for the subsequence $\{N''\}$ of multiples of q.

The proof in this case is complete.

6. The case where $0 \leq r < 1/6$.

In this section we will show the following proposition.

Proposition 6.1. *For any r with $0 \leq r < 1/6$, we have*

$$\lim_{N\to\infty} \frac{\log|J_N(E;\exp(2\pi r\sqrt{-1}/N))|}{N} = 0.$$

Proof. Since $J_N(E;1) = 1$, we have the equality when $r = 0$.

If $0 < r < 1/6$, then $-1 < g(j) < 0$ for $j = 1, 2, \ldots, N-1$ and so $|f(k)| < 1$ for $k = 1, 2, \ldots, N-1$. Therefore

$$\left|J_N\left(E;\exp(2\pi r\sqrt{-1}/N)\right)\right| = \left|\sum_{k=0}^{N-1} f(k)\right| < N.$$

On the other hand, since $f(k)$ is positive (negative, respectively) if k is even (odd, respectively) and $|f(k)|$ is decreasing, we have

$$\left|\sum_{k=0}^{N-1} f(k)\right|$$

$$
=\begin{cases}
\{1+f(1)\}+\{f(2)+f(3)\}+\cdots+\{f(N-2)+f(N-1)\} \\
\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{if } N \text{ is even} \\
\{1+f(1)\}+\{f(2)+f(3)\}+\cdots+\{f(N-3)+f(N-2)\}+f(N-1) \\
\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \text{if } N \text{ is odd}
\end{cases}
$$

$$
> 1+f(1) = 1+2\cos(2\pi r)-2\cos(2\pi r/N) > (2\pi r)^6/360N^6.
$$

Therefore we have

$$
\frac{\log\left((2\pi r)^6/360N^6\right)}{N} < \frac{\log\left(J_N\left(E;\exp(2\pi r\sqrt{-1}/N)\right)\right)}{N} < \frac{\log N}{N}
$$

and the proposition follows. □

7. The case where r is irrational and $1/6 \le |1-r| < 1/4$.

In this section we will prove a result similar to Theorem1.2 for irrational r with $1/6 \le |1-r| < 1/4$.

Proposition 7.1. *If r is irrational and $1/6 \le |1-r| < 1/4$, then we have*

$$
\limsup_{N\to\infty} 2\pi\frac{\log\left|J_N\left(E;\exp(2\pi r\sqrt{-1}/N)\right)\right|}{N} = \frac{2\Lambda\bigl(\pi r+\theta(r)/2\bigr)-2\Lambda\bigl(\pi r-\theta(r)/2\bigr)}{r}.
$$

First we prove the proposition in the case where $3/4 < r \le 5/6$.

Proof when $3/4 < r \le 5/6$. We put

$$
A := \frac{N\varphi(r)}{2\pi r}, \quad B := \frac{N(1-r)}{r}, \quad C := \frac{N\theta(r)}{2\pi r}, \quad \text{and} \quad D := \frac{N(2\pi-\theta(r))}{2\pi r},
$$

with $\varphi(r) := \arccos(\cos(2\pi r)+1/2)$ and $\theta(r) := \arccos(\cos(2\pi r)-1/2)$. Note that $g(A)=-1$, $g(B)=0$, and $g(C)=g(D)=1$.

Since

$$
\begin{aligned}
g(j) &< -1 &&\text{for} && j < A, \\
-1 < g(j) &< 0 &&\text{for} && A < j < B, \\
0 < g(j) &< 1 &&\text{for} && B < j < C, \\
g(j) &> 1 &&\text{for} && C < j < D, \quad \text{and} \\
0 < g(j) &< 1 &&\text{for} && D < j,
\end{aligned}
$$

Table of $g(j)$ regarding j as a continuous parameter

j	0	$\cdots$	A	$\cdots$	B	$\cdots$	C	$\cdots$	D	$\cdots$	N
$g(j)$	$g(0) < -1$	↗	-1	↗	0	↗	1	$g(j) > 1$	1	↘	0

we have

- $1 = |f(0)| < |f(1)| < \cdots < |f(\lfloor A\rfloor - 1)| < |f(\lfloor A\rfloor)|$,
- $|f(\lfloor A\rfloor)| > |f(\lfloor A\rfloor+1)| > \cdots > |f(\lfloor B\rfloor-1)| > |f(\lfloor B\rfloor)| > |f(\lfloor B\rfloor+1)| > \cdots > |f(\lfloor C\rfloor-1)| > |f(\lfloor C\rfloor)|$,
- $|f(\lfloor C\rfloor)| < |f(\lfloor C\rfloor+1)| < \cdots < |f(\lfloor D\rfloor-1)| < |f(\lfloor D\rfloor)|$, and
- $|f(\lfloor D\rfloor)| > |f(\lfloor D\rfloor+1)| > \cdots > |f(N-2)| > |f(N-1)|$.

Therefore $|f(k)|$ takes its 'local maxima' at $k = \lfloor A\rfloor$ and $\lfloor D\rfloor$.

Now we have

$$
\begin{aligned}
&\lim_{N\to\infty}\frac{\log|f(\lfloor A\rfloor)|}{N} \\
=&\lim_{N\to\infty}\frac{1}{N}\sum_{j=1}^{\lfloor A\rfloor}\{\log|2\sin(\pi rj/N+\pi r)| + \log|2\sin(\pi rj/N-\pi r)|\} \\
=&\frac{1}{\pi r}\int_{\pi r}^{\varphi(r)/2+\pi r}\log|2\sin y|\,dy + \frac{1}{\pi r}\int_{-\pi r}^{\varphi(r)/2-\pi r}\log|2\sin y|\,dy
\end{aligned}
$$

$$=\frac{\Lambda(\pi r-\varphi(r)/2)-\Lambda(\pi r+\varphi(r)/2)}{\pi r}$$

and similarly

$$\begin{aligned}&\lim_{N\to\infty}\frac{\log|f(\lfloor D\rfloor)|}{N}\\ =&\frac{1}{\pi r}\int_{\pi r}^{\pi-\theta(r)/2+\pi r}\log|2\sin y|\,dy+\frac{1}{\pi r}\int_{-\pi r}^{\pi-\theta(r)/2-\pi r}\log|2\sin y|\,dy\\ =&\frac{\Lambda(\pi r+\theta(r)/2)-\Lambda(\pi r-\theta(r)/2)}{\pi r}.\end{aligned}$$

Since

$$\begin{aligned}(7.1)\quad \delta:=&\pi r\left(\lim_{N\to\infty}\frac{\log|f(\lfloor D\rfloor)|}{N}-\lim_{N\to\infty}\frac{\log|f(\lfloor A\rfloor)|}{N}\right)\\ =&\Lambda(\pi r+\theta(r)/2)-\Lambda(\pi r-\theta(r)/2)+\Lambda(\pi r+\varphi(r)/2)-\Lambda(\pi r-\varphi(r)/2)>0,\end{aligned}$$

(see Appendix) we see that $|f(\lfloor D\rfloor)|$ is the maximum, that is, $|f(k)|\le|f(\lfloor D\rfloor)|$ for sufficient large N. Therefore $|J_N(E;\exp(2\pi r\sqrt{-1}/N))|<N|f(\lfloor D\rfloor)|$.

On the other hand, since the sign of $f(k)$ does not change for $k\ge\lfloor B\rfloor$ and $|f(k)|<|f(\lfloor A\rfloor)|$ for $k\le\lfloor C\rfloor$, we have

$$\begin{aligned}&|J_N(E;\exp(2\pi r\sqrt{-1}/N))|\\ &\qquad=\left|\sum_{k=0}^{N-1}f(k)\right|>\left|f(\lfloor D\rfloor)-\sum_{k=0}^{\lfloor B-1\rfloor}f(k)\right|>|f(\lfloor D\rfloor)-Nf(\lfloor A\rfloor)|.\end{aligned}$$

Since

$$\frac{|f(\lfloor A\rfloor)|}{|f(\lfloor D\rfloor)|}\underset{N\to\infty}{\sim}\exp(-\delta N)$$

from (7.1), we have

$$\begin{aligned}|J_N(E;\exp(2\pi r\sqrt{-1}/N))|>&|f(\lfloor D\rfloor)|\left(1-N\frac{f(\lfloor A\rfloor)}{f(\lfloor D\rfloor)}\right)\\ &\underset{N\to\infty}{\sim}|f(\lfloor D\rfloor)|\{1-N\exp(-\delta N)\}\\ &\xrightarrow{N\to\infty}|f(\lfloor D\rfloor)|.\end{aligned}$$

So we finally have

$$\begin{aligned}\lim_{N\to\infty}\frac{\log|J_N(E;\exp(2\pi r\sqrt{-1}/N))|}{N}=&\lim_{N\to\infty}\frac{\log|f(\lfloor D\rfloor)|}{N}\\ =&\frac{\Lambda(\pi r+\theta(r)/2)-\Lambda(\pi r-\theta(r)/2)}{\pi r}\end{aligned}$$

as required. □

Next we will consider the case where $7/6\le r<5/4$.

Proof when $7/6\le r<5/4$. We put

$$A:=\frac{N\varphi(r)}{2\pi r},\quad B:=\frac{N(r-1)}{r},\quad C:=\frac{N\theta(r)}{2\pi r},\quad D:=\frac{N(2\pi-\theta(r))}{2\pi r},$$
$$B':=\frac{N(2-r)}{r},\quad A':=\frac{N(2\pi-\varphi(r))}{r},\quad\text{and}\quad A'':=\frac{N(2\pi+\varphi(r))}{r}$$

with $\varphi(r):=\arccos(\cos(2\pi r)+1/2)$ and Note that $g(A)=g(A')=g(A'')=-1$, $g(B)=g(B')=0$, and $g(C)=g(D)=1$.

Since

$$\begin{aligned} g(j) &< -1 && \text{for} && j < A, \\ -1 < g(j) &< 0 && \text{for} && A < j < B, \\ 0 < g(j) &< 1 && \text{for} && B < j < C, \\ g(j) &> 1 && \text{for} && C < j < D, \\ 0 < g(j) &< 1 && \text{for} && D < j < B', \\ -1 < g(j) &< 0 && \text{for} && B' < j < A', \\ g(j) &< -1 && \text{for} && A' < j < A'', \quad \text{and} \\ -1 < g(j) &< 0 && \text{for} && j > A'', \end{aligned}$$

$|f(k)|$ has three 'local maxima' at $k = \lfloor A \rfloor$, $\lfloor D \rfloor$, and $\lfloor A'' \rfloor$.

But since

$$\begin{aligned} &\lim_{N\to\infty} \frac{\log |f(\lfloor A'' \rfloor)|}{N} \\ =&\lim_{N\to\infty} \frac{1}{N} \sum_{j=1}^{\lfloor A'' \rfloor} \{\log|2\sin(\pi r j/N + \pi r)| + \log|2\sin(\pi r j/N - \pi r)|\} \\ =&\frac{1}{\pi r}\int_{\pi r}^{\pi+\varphi(r)/2+\pi r} \log|2\sin y|\,dy + \frac{1}{\pi r}\int_{-\pi r}^{\pi+\varphi(r)/2-\pi r} \log|2\sin y|\,dy \\ =&\frac{\Lambda(\pi r - \varphi(r)/2) - \Lambda(\pi r + \varphi(r)/2)}{\pi r} < 0, \end{aligned}$$

$|f(\lfloor D \rfloor)|$ is the unique maximum. Therefore the argument for the case $3/4 < r \le 5/6$ can be applied and the proof is complete. $\square$

8. The case where r is purely imaginary.

In this section we consider the case where $t = \exp(2\pi r/N)$.

Theorem 8.1. *If* $2\pi|r| > \operatorname{arccosh}(3/2)$*, then*

$$2\pi|r| \lim_{N\to\infty} \frac{\log J_N(E;\exp(2\pi r/N))}{N} = 2\Gamma(\pi|r| + \varphi(|r|)/2) - 2\Gamma(\pi|r| - \varphi(|r|)/2),$$

where $\Gamma(z) := \int_0^z \log 2\sinh x d x$ *and* $\varphi(r) := \operatorname{arccosh}(\cosh(2\pi r) - 1/2)$*. If* $2\pi|r| \le \operatorname{arccosh}(3/2)$*, then*

$$\lim_{N\to\infty} \frac{\log J_N(E;\exp(2\pi r/N))}{N} = 0.$$

Proof. In this case

$$\begin{aligned} g(j) &= 4\sinh(\pi|r|j/N + \pi|r|)\sinh(\pi|r|j/N - \pi|r|) \\ &= 2\cosh(2\pi|r|) - 2\cosh(2\pi|r|j/N) > 0. \end{aligned}$$

Since $g(j)$ is monotonically decreasing and $g(0) = 2\cosh(2\pi|r|) - 2$, $f(k) = \prod_{j=1}^{k} g(j)$ takes its maximum at $g(\lfloor \varphi(|r|)N/2\pi|r| \rfloor)$ if $g(0) > 1$. Therefore

$$\begin{aligned} &\lim_{N\to\infty} \frac{\log J_N E;\exp(2\pi|r|/N)}{N} \\ &= \lim_{N\to\infty} \frac{\log f(\lfloor \varphi(|r|)N/2\pi|r| \rfloor)}{N} \\ &= \lim_{N\to\infty} \sum_{j=1}^{\lfloor \varphi(|r|)N/2\pi|r| \rfloor} \frac{\log 2\sinh(\pi|r|j/N + \pi|r|)}{N} \\ &\quad + \lim_{N\to\infty} \sum_{j=1}^{\lfloor \varphi(|r|)N/2\pi|r| \rfloor} \frac{\log 2\sinh(\pi|r|j/N - \pi|r|)}{N} \end{aligned}$$

$$= \frac{1}{\pi|r|}\int_{\pi|r|}^{\varphi(|r|)/2+\pi|r|}\log 2\sinh x\,d\,x + \frac{1}{\pi|r|}\int_{-\pi|r|}^{\varphi(|r|)/2-\pi|r|}\log 2\sinh x\,d\,x$$

and the result follows when $2\pi|r| > \operatorname{arccosh}(3/2)$.

If $2\pi|r| \le \operatorname{arccosh}(3/2)$, then $g(j) < g(0) = 2\cosh(2\pi|r|) - 2 \le 1$ and so

$$N > J_N(J; \exp(2\pi|r|/N)) > g(0) - \varepsilon = 2\cosh(2\pi r) - 2 - \varepsilon$$

for any $\varepsilon > 0$ if N is sufficiently large. Therefore

$$\lim_{N\to\infty}\frac{\log J_N(E; \exp(2\pi r/N))}{N} = 0$$

as required. □

Acknowledgments. Part of this work was done when the author was visiting the International Institute for Advanced Studies from 5th to 8th March, 2002 to attend the workshop 'Volume Conjecture and Its Related Topics' (partially supported by the Research Institute for Mathematical Sciences, Kyoto University), Warwick University to attend the workshop 'Quantum Topology' from 18th to 22nd March, 2002, and Université du Québec à Montréal to attend the workshop 'Knots in Montreal II' from 20th to 21st April, 2002. Thanks are due to the institute/universities and to the organizers of the workshops, Stavros Garoufalidis, Colin Rourke, Steven Boyer, and Adam Sikora.

The author thanks K. Hikami for introducing the computer program PARI. He is also grateful to S. Gukov, A. Mednykh, and A. Vesnin for informing their results.

Appendix A. Some calculations

Lemma A.1. *For* $|r-1| < 1/3$*, we have*

$$\Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2) > 0$$

Proof. Put $V(r) := \Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2)$ Since

$$4\sin(\pi r + \theta(r)/2)\sin(\pi r - \theta(r)/2) = 2\cos(\theta(r)) - 2\cos(2\pi r) = -1,$$

we have

$$\begin{aligned}\frac{d\,V(r)}{d\,r} &= -\log|2\sin(\pi r + \theta(r)/2)|\left(\pi + \frac{1}{2}\frac{d\,\theta(r)}{d\,r}\right)\\ &\quad + \log|2\sin(\pi r - \theta(r)/2)|\left(\pi - \frac{1}{2}\frac{d\,\theta(r)}{d\,r}\right)\\ &= 2\pi\log|2\sin(\pi r - \theta(r)/2)|\,.\end{aligned}$$

Since $\pi/6 \le \pi r - \theta(r)/2 < 5\pi/6$ for $2/3 < r < 1$, $\sin(\pi r - \theta(r)/2) > 1/2$ and so $d\,V(r)/d\,r > 0$. Therefore $V(r) \ge 0$ for $2/3 < r < 1$ since $V(2/3) = 0$.

Since $V(2-r) = V(r)$, the equality also holds for $1 < r < 4/3$. □

Lemma A.2. *For* $1/6 < |r-1| < 1/4$*, we have*

$$\Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2) + \Lambda(\pi r + \varphi(r)/2) - \Lambda(\pi r - \varphi(r)/2) > 0$$

Proof. Put $W(r) := \Lambda(\pi r + \theta(r)/2) - \Lambda(\pi r - \theta(r)/2) + \Lambda(\pi r + \varphi(r)/2) - \Lambda(\pi r - \varphi(r)/2)$. As in the proof of the previous lemma, we have

$$\frac{d\,W(r)}{d\,r} = 2\log|2\sin(\pi r - \theta(r)/2)| + 2\log|2\sin(\pi r - \varphi(r)/2)|.$$

Since $7\pi/12 < \pi r - \varphi(r)/2 < 5\pi/6$ for $3/4 < r < 5/6$, $\sin(\pi r - \varphi(r)/2) > 1/2$ and so $d\,W(r)/d\,r > 0$. Therefore $W(r) > 0$ for $3/4 < r < 5/6$ since $V(3/4) = 0$.

Since $W(2-r) = W(r)$, the equality also holds for $7/6 < r < 5/4$. □

References

[1] S. Gukov, *Three-Dimensional Quantum Gravity, Chern-Simons Theory, and the A-Polynomial*, HUTP-03/A003, ITEP-TH-50/02, arXiv:hep-th/0306165.

[2] R. M. Kashaev, *A link invariant from quantum dilogarithm*, Modern Phys. Lett. A **10** (1995), no. 19, 1409–1418. MR **96j:**81060

[3] ———, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. **39** (1997), no. 3, 269–275. MR **98b:**57012

[4] R. M. Kashaev and O. Tirkkonen, *A proof of the volume conjecture on torus knots*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **269** (2000), no. Vopr. Kvant. Teor. Polya i Stat. Fiz. 16, 262–268, 370. MR 1 805 865

[5] H. Murakami, *The asymptotic behavior of the colored Jones function of a knot and its volume*, Proceedings of 'Art of Low Dimensional Topology VI' (T. Kohno, ed.), January 2000, arXiv:math.GT/0004036.

[6] ———, *Optimistic calculations about the Witten-Reshetikhin-Turaev invariants of closed three-manifolds obtained from the figure-eight knot by integral Dehn surgeries*, Sūrikaisekikenkyūsho Kōkyūroku (2000), no. 1172, 70–79. MR 1 805 729

[7] ———, *Kashaev's invariant and the volume of a hyperbolic knot after Y. Yokota*, Physics and combinatorics 1999 (Nagoya), World Sci. Publishing, River Edge, NJ, 2001, pp. 244–272. MR **2002k:**57031

[8] ———, *Mahler measure of the colored Jones polynomial and the volume conjecture*, Sūrikaisekikenkyūsho Kōkyūroku (2002), no. 1279, 86–99. MR 1 953 834

[9] H. Murakami and J. Murakami, *The colored Jones polynomials and the simplicial volume of a knot*, Acta Math. **186** (2001), no. 1, 85–104. MR 1 828 373

[10] H. Murakami, J. Murakami, M. Okamoto, T. Takata, and Y. Yokota, *Kashaev's conjecture and the Chern-Simons invariants of knots and links*, Experiment. Math. **11** (2002), no. 3, 427–435. MR 1 959 752

[11] A. Yu. Vesnin and A. D. Mednykh, *Hyperbolic volumes of Fibonacci manifolds*, Sibirsk. Mat. Zh. **36** (1995), no. 2, 266–277, i. MR **97c:**57019

[12] Y. Yokota, *On the volume conjecture for hyperbolic knots*, arXiv:math.QA/0009165.

[13] ———, *On the volume conjecture for hyperbolic knots*, Proceedings of the 47th Topology Symposium (Inamori Hall, Kagoshima University), July 2000, pp. 38–44.

[14] ———, *On the volume conjecture of hyperbolic knots*, Knot Theory – dedicated to Professor Kunio Murasugi for his 70th birthday (M. Sakuma, ed.), March 2000, pp. 362–367.

[15] ———, *On the potential functions for the hyperbolic structures of a knot complement*, Geom. Topol. Monogr. **4** (2002), 303–311.

[16] ———, *From the Jones polynomial to the A-polynomial of hyperbolic knots*, Interdiscip. Inform. Sci. **9** (2003), 11–21.

Department of Mathematics, Tokyo Institute of Technology, Oh-okayama, Meguro, Tokyo 152-8551, Japan

E-mail address: `starshea@tky3.3web.ne.jp`